
Huff Creek














This doc

may not be rec

duced in whole or in part without the express written consent of NAI Columbi



Map Updated: Friday, Se

ber 15, 2023. This

mitted is not guaranteed. Although obtained from reliable sources, all inf

SPECIAL FLOOD HAZARD AREAS

1% Annual Chance Flood Hazard Zone A. A.E. A98.A.O. A.H. A.R. K. VE

Regulatory Floodway

0THER AREAS OF FLOOD HAZARD 0.2% Annua I Chance Flood Hazard Zoos %

Zone X Future Conditions 1% Annual Chance Flood Hazard Zone X Area with Reduced Flood Risk due to Levee Zone X MO SCREEN Areas Outside the 0.2% Annual Chance Floodplain Zone X Areas of Undetermined Flood Hazard Zone D

CROSS SECTIONS & BFES

E) 17.5	Cross Sections with 1% Annual Chance Water Surface Elevation
6)	Coastal Transect
	Coastal Transect Baseline
	Profile Baseline
	Base Flood Elevation

rossing Map Updated: Friday, September 15, 2023, This ces, all info

1165

n should be confir

or to use or reliance upon the


0


Hope






National Wetlands Inventory











Map Unit Description (Brief, Generated)

Pickens County, South Carolina

[Minor map unit components are excluded from this report]

Map unit: PaC2 - Pacolet fine sandy loam, 6 to 10 percent slopes, eroded

Component: Pacolet (100%)

The Pacolet component makes up 100 percent of the map unit. Slopes are 6 to 10 percent. This component is on hillslopes on piedmonts. The parent material consists of clayey residuum weathered from granite and gneiss. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is well drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches is moderate. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 3e. This soil does not meet hydric criteria.

Map unit: PaE2 - Pacolet fine sandy loam, 10 to 25 percent slopes, eroded

Component: Pacolet (100%)

The Pacolet component makes up 100 percent of the map unit. Slopes are 10 to 25 percent. This component is on hillslopes on piedmonts. The parent material consists of clayey residuum weathered from granite and gneiss. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is well drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches is moderate. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 6e. This soil does not meet hydric criteria.

Map unit: PaF - Pacolet fine sandy loam, 25 to 40 percent slopes

Component: Pacolet (100%)

The Pacolet component makes up 100 percent of the map unit. Slopes are 25 to 40 percent. This component is on hillslopes on piedmonts. The parent material consists of clayey residuum weathered from granite and gneiss. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is well drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches is moderate. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 7e. This soil does not meet hydric criteria.

Map unit: PcC3 - Pacolet clay loam, 2 to 10 percent slopes, severely eroded

Component: Pacolet, severely eroded (100%)

The Pacolet component makes up 100 percent of the map unit. Slopes are 2 to 10 percent. This component is on hillslopes on piedmonts. The parent material consists of clayey residuum weathered from granite and gneiss. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is well drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches is moderate. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 4e. This soil does not meet hydric criteria.



[Minor map unit components are excluded from this report]

Map unit: PcE3 - Pacolet clay loam, 10 to 25 percent slopes, severely eroded

Component: Pacolet, severely eroded (100%)

The Pacolet component makes up 100 percent of the map unit. Slopes are 10 to 25 percent. This component is on hillslopes on piedmonts. The parent material consists of clavey residuum weathered from granite and gneiss. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is well drained. Water movement in the most restrictive layer is moderately high. Available water to a depth of 60 inches is moderate. Shrink-swell potential is low. This soil is not flooded. It is not ponded. There is no zone of water saturation within a depth of 72 inches. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 7e. This soil does not meet hydric criteria.

Map unit: To - Toccoa soils

Component: Toccoa (100%)

The Toccoa component makes up 100 percent of the map unit. Slopes are 0 to 2 percent. This component is on flood plains on piedmonts. The parent material consists of loamy alluvium. Depth to a root restrictive layer is greater than 60 inches. The natural drainage class is moderately well drained. Water movement in the most restrictive layer is high. Available water to a depth of 60 inches is moderate. Shrink-swell potential is low. This soil is occasionally flooded. It is not ponded. A seasonal zone of water saturation is at 45 inches during January, February, March, April, December. Organic matter content in the surface horizon is about 1 percent. Nonirrigated land capability classification is 2w. This soil does not meet hydric criteria.

Map Unit Description (Brief, Generated)



